

DR-59. THE REACTION 5-HETARYL(ARYL)METHYLIDENE-2,4,6-PYRIMIDINE-2,4,6(1H,3H,5H)-TRIONS WITH *L*-PROLINE AND PARAFORMES


A. G. Tyrkov¹, S. A. Luzhnova², E. A. Yurtaeva²

¹ Astrakhan State University, Tatischev St., 20a, Astrakhan, 414000, Russia

² Pyatigorsk Medical Pharmaceutical Institute, Kalinin Av., 11, Pyatigorsk, 357532, Russia

E-mail: tyrkov@rambler.ru

In the literature, there are limited information on the use of azomethynyls in the reactions of 1,3-dipolar cycloaddition [1, 2]. We found that the reaction of the three-component heterocyclization of pyrimidine-2,4,6-trions **1–3** with *L*-proline and paraffin proceeds with heating for 17 hours in dried toluene and yields, in 80–82 % yield, the previously unknown spirocompounds **4–6**.



Het = бензофуран-2-ил (**1**); 1,3-бензотиозол-2-ил (**2**); 2,1,3-бензооксадиазол-5-ил (**3**)

It can be assumed that during the thermolysis of paraform with *L*-proline, corresponding azomethonilides **A** are generated, the latter as a result of the 1,3-dipolar cycloaddition process to the molecules of dipolarophilic compounds **1–3** stabilize to the target compounds **4–6**.

References

1. Furan-related N-onium salts as synthons for preparation of 3-furylbenzoindolizines / J. Štetinová [et al.] // Collect. Czechoslov. Chem. Commun. Institute of Organic Chemistry and Biochemistry AS CR, v.v.i., 1986. Vol. 51, № 2. P. 412.
2. Reactions of 2-aryl-1,1-dicyanoethenes with *L*-proline and aldehydes / S. B. Nosachev [et al.] // Russ. J. Org. Chem. SP MAIK Nauka / Interperiodica. 2010. Vol. 46, № 5. P. 674.